

Computational Geometry**Exercise Set 5****HS07**URL: <http://www.ti.inf.ethz.ch/ew/courses/CG07/>**Exercise 1 (10 points)**

You are given

- a star-shaped polygon $P \subset \mathbb{R}^2$, represented as a doubly connected list of its vertices $V(P)$,
- and a point $c \in P$ (not necessarily in $V(P)$), such that for all $p \in P$ the line segment \overline{cp} is contained in P .

Describe an algorithm which triangulates P in linear time. The algorithm could for example output all edges of the triangulation, that are not already edges of the polygon.

Exercise 2 (10 points)

This exercise is about an application from *Computational Biology*:

You are given a set of disks $P = \{a_1, \dots, a_n\}$ in \mathbb{R}^2 , all with the same radius $r_a > 0$. Each of these disks represents an atom of a protein. A water molecule is represented by a disc with radius $r_w > r_a$. A water molecule cannot intersect the interior of any protein atom, but it can be tangent to one. We say that an atom $a_i \in P$ is *solvent-accessible* if there exists a placement of a water molecule such that it is tangent to a_i and does not intersect the interior of any other atom in P . Given P , find an $O(n \log n)$ time algorithm which determines all solvent-inaccessible molecules of P .


Exercise 3 (10 points)

Consider the unit paraboloid $\mathcal{U} : z = x^2 + y^2$ in \mathbb{R}^3 and let $u : p = (p_x, p_y, 0) \mapsto (p_x, p_y, p_x^2 + p_y^2)$ be the orthogonal projection of the x/y -plane onto \mathcal{U} . What is the equation for the tangent plane H_p to \mathcal{U} in $u(p)$?

Let p and q be two points in the x/y -plane and $h_p : \mathbb{R}^3 \rightarrow H_p$ the orthogonal projection (i.e. in z -direction) of the x/y -plane onto H_p . Show:

$$\|u(q) - h_p(q)\| = \|p - q\|^2 .$$

Here is an illustration:



Due date: 5.11.2007