

## RECAP: Extremal problems — Examples

**Proposition 1.** If  $G$  is an  $n$ -vertex graph with **at most**  $n - 2$  edges then  $G$  is disconnected.

**A Question** you always have to ask:

Can we improve on this proposition?

**Answer.** NO! The same statement is **FALSE** with  $n - 1$  in the place of  $n - 2$ .

Proposition 1 is **best possible**, as shown by  $P_n$ .

**Proposition 1. +  $P_n$ :** The **minimum** value of  $e(G)$  over connected graphs is  $n - 1$ .

**Proposition 2.** If  $G$  is an  $n$ -vertex graph with **at least**  $n$  edges then  $G$  contains a cycle.

**Remark.** Proposition 2 is also **best possible**, (e.g.  $P_n$ ).

**Proposition 2. + Remark:** The **maximum** value of  $e(G)$  over acyclic (i.e. cycle-free) graphs is  $n - 1$ .

## Extremal problems — More example\_\_\_\_\_

Vague description: An **extremal problem** asks for the maximum or minimum value of a parameter over a class of objects (graphs, in most cases).

**Proposition.**  $G$  is an  $n$ -vertex graph with  $\delta(G) \geq \lfloor n/2 \rfloor$ , then  $G$  is connected.

**Remark.** The above proposition is **best possible**, as shown by  $K_{\lfloor n/2 \rfloor} + K_{\lceil n/2 \rceil}$ .

Graph  $G + H$  is the **disjoint union** (or **sum**) of graphs  $G$  and  $H$ . For an integer  $m$ ,  $mG$  is the graph consisting of  $m$  disjoint copies of  $G$ .

**Prop. + Remark:** The **maximum** value of  $\delta(G)$  over disconnected graphs is  $\lfloor \frac{n}{2} \rfloor - 1$ .

# Extremal Problems\_\_\_\_\_

graph property	graph parameter	type of extremum	value of extremum
connected	$e(G)$	minimum	$n - 1$
acyclic	$e(G)$	maximum	$n - 1$
disconnected	$\delta(G)$	maximum	$\lfloor \frac{n}{2} \rfloor - 1$
$K_3$ -free	$e(G)$	maximum	$\lfloor \frac{n^2}{4} \rfloor$

## Triangle-free subgraphs\_\_\_\_\_

**Theorem.** (Mantel, 1907) The maximum number of edges in an  $n$ -vertex **triangle-free** graph is  $\lfloor \frac{n^2}{4} \rfloor$ .

*Proof.*

(i) *There is a triangle-free graph with  $\lfloor \frac{n^2}{4} \rfloor$  edges.*

(ii) *If  $G$  is a triangle-free graph, then  $e(G) \leq \lfloor \frac{n^2}{4} \rfloor$ .*

Proof of (ii) is with extremality. (Look at the neighborhood of a vertex of maximum degree.)

## Complete $k$ -partite graphs\_\_\_\_\_

$G$  is a **complete  $k$ -partite graph** if there is a partition  $V_1 \cup \dots \cup V_k = V(G)$  of the vertex set, such that  $uv \in E(G)$  iff  $u$  and  $v$  are in *different* parts of the partition. If  $|V_i| = n_i$ , then  $G$  is denoted by  $K_{n_1, \dots, n_k}$ .

The **Turán graph  $T_{n,r}$**  is the complete  $r$ -partite graph on  $n$  vertices whose partite sets differ in size by at most 1. (All partite sets have size  $\lceil n/r \rceil$  or  $\lfloor n/r \rfloor$ .)

**Lemma** Among  $r$ -colorable graphs the Turán graph is the *unique* graph, which has the most number of edges.

*Proof.* Local change. □

## Turán's Theorem

---

The **Turán number**  $ex(n, H)$  of a graph  $H$  is the largest integer  $m$  such that there exists an  $H$ -free\* graph on  $n$  vertices with  $m$  edges.

*Example:* Mantel's Theorem states  $ex(n, K_3) = \lfloor \frac{n^2}{4} \rfloor$ .

**Theorem.** (Turán, 1941)

$$ex(n, K_r) = e(T_{n,r-1}) = \left(1 - \frac{1}{r-1}\right) \binom{n}{2} + O(n).$$

*Proof.* Prove by induction on  $r$  that


$G \not\supseteq K_r \implies$  **there is** an  $(r-1)$ -partite graph  $H$  with  $V(H) = V(G)$  and  $e(H) \geq e(G)$ .

Then apply the Lemma to finish the proof. □

\*Here  $H$ -free means that there is no subgraph isomorphic to  $H$

## Turán-type problems\_\_\_\_\_

**Question.** (Turán, 1941) What happens if instead of  $K_4$ , which is the graph of the tetrahedron, we forbid the graph of some other platonic polyhedra? How many edges can a graph without an octahedron (or cube, or dodecahedron or icosahedron) have?



The platonic solids

## Erdős-Simonovits-Stone Theorem\_\_\_\_\_

**Theorem.** (Erdős-Stone, 1946) For arbitrary fixed integers  $r \geq 2$  and  $t \geq 1$

$$ex(n, T_{rt,r}) = \left(1 - \frac{1}{r-1}\right) \binom{n}{2} + o(n^2).$$

**Corollary.** (Erdős-Simonovits, 1966) For any graph  $H$ ,

$$ex(n, H) = \left(1 - \frac{1}{\chi(H)-1}\right) \binom{n}{2} + o(n^2).$$

### Corollaries of the Corollary.

$$\begin{aligned} ex(n, \text{octahedron}) &= \frac{n^2}{4} + o(n^2) \\ ex(n, \text{dodecahedron}) &= \frac{n^2}{4} + o(n^2) \\ ex(n, \text{icosahedron}) &= \frac{n^2}{3} + o(n^2) \\ ex(n, \text{cube}) &= o(n^2) \end{aligned}$$



## Proof of the Erdős-Simonovits Corollary\_\_\_\_\_

**Theorem.** (Erdős-Stone, 1946) For arbitrary fixed integers  $r \geq 2$  and  $t \geq 1$

$$ex(n, T_{rt,r}) = \left(1 - \frac{1}{r-1}\right) \binom{n}{2} + o(n^2).$$

**Corollary.** (Erdős-Simonovits, 1966) For any graph  $H$ ,

$$ex(n, H) = \left(1 - \frac{1}{\chi(H)-1}\right) \binom{n}{2} + o(n^2).$$

*Proof of the Corollary.* Let  $r = \chi(H)$ .

- $\chi(T_{n,r-1}) < \chi(H)$ , so  $e(T_{n,r-1}) \leq ex(n, H)$ .
- $T_{r\alpha,r} \supseteq H$ , so  $ex(n, T_{r\alpha,r}) \geq ex(n, H)$ , where  $\alpha$  is a constant depending on  $H$ ; say  $\alpha = \alpha(H)$ .

□

The number of edges in a  $C_4$ -free graph\_\_\_\_\_

**Theorem** (Erdős, 1938)  $ex(n, C_4) = O(n^{3/2})$

*Proof.* Let  $G$  be a  $C_4$ -free graph on  $n$  vertices.

$C = C(G) :=$  number of  $K_{1,2}$  (“cherries”) in  $G$ .

*Doublecount*  $C$ .

**Counting by the midpoint:** Every vertex  $v$  is the midpoint of exactly  $\binom{d(v)}{2}$  cherries. Hence

$$C = \sum_{v \in V} \binom{d(v)}{2}.$$

**Counting by the endpoints:** Every pair  $\{u, w\}$  of vertices form the endpoints of **at most one** cherry. (Otherwise there is a  $C_4 \subseteq G$ .) Hence

$$C \leq 1 \cdot \binom{n}{2}.$$

## Proof cont'd

---

Combine and apply Jensen's inequality  
(Note that  $x \rightarrow \binom{x}{2}$  is a convex function)

$$\binom{n}{2} \geq C \geq \sum_{v \in V} \binom{d(v)}{2} \geq n \cdot \binom{\bar{d}(G)}{2}.$$

$\bar{d}(G) = \frac{1}{n} \sum_{v \in V} d(v)$  is the **average degree** of  $G$ .

$$\frac{n-1}{2} \geq \binom{\bar{d}(G)}{2} \geq \frac{(\bar{d}(G)-1)^2}{2}$$

Hence  $\sqrt{n-1} + 1 \geq \bar{d}(G)$ . □

**Theorem** (E. Klein, 1938)  $ex(n, C_4) = \Theta(n^{3/2})$

*Proof.* Homework.

**Theorem** (Kővári-Sós-Turán, 1954) For  $s \geq t \geq 1$

$$ex(n, K_{t,s}) \leq c_s n^{2-\frac{1}{t}}$$

*Proof.* Homework.

# Open problems and Conjectures\_\_\_\_\_

## Known results.

$$\Omega(n^{3/2}) \leq ex(n, Q_3) \leq O(n^{8/5})$$

$$\Omega(n^{9/8}) \leq ex(n, C_8) \leq O(n^{5/4})$$

$$\Omega(n^{5/3}) \leq ex(n, K_{4,4}) \leq O(n^{7/4})$$

## Conjectures.

$$ex(n, K_{t,s}) = \Theta\left(n^{2 - \frac{1}{\min\{t,s\}}}\right) \text{ true for } t = 2, 3 \text{ and } s \geq t \\ \text{or } t \geq 4 \text{ and } s > (t - 1)!$$

$$ex(n, C_{2k}) = \Theta\left(n^{1 + \frac{1}{k}}\right) \text{ true for } k = 2, 3 \text{ and } 5$$

$$ex(n, Q_3) = \Theta\left(n^{\frac{8}{5}}\right)$$

If  $H$  is a  $d$ -degenerate bipartite graph, then

$$ex(n, H) = O\left(n^{2 - \frac{1}{d}}\right).$$