

Extremal problems* — Examples_____

Proposition. G is an n -vertex graph with $\delta(G) \geq \lfloor n/2 \rfloor$, then G is connected.

Remark. The above proposition is *best possible*, as shown by $K_{\lfloor n/2 \rfloor} + K_{\lceil n/2 \rceil}$.

Graph $G + H$ is the **disjoint union** (or **sum**) of graphs G and H . For an integer m , mG is the graph consisting of m disjoint copies of G .

Prop. + Remark: The **maximum** value of $\delta(G)$ over disconnected graphs is $\lfloor \frac{n}{2} \rfloor - 1$.

Vague description: An **extremal problem** asks for the maximum or minimum value of a parameter over a class of objects (graphs, in most cases).

Proposition. The **minimum** number of edges in a connected graph is $n - 1$.

*My favorite topic

Extremal Problems_____

graph property	graph parameter	type of extremum	value of extremum
connected	$e(G)$	minimum	$n - 1$
disconnected	$\delta(G)$	maximum	$\lfloor \frac{n}{2} \rfloor - 1$
K_3 -free	$e(G)$	maximum	$\lfloor \frac{n^2}{4} \rfloor$

Triangle-free subgraphs_____

Theorem. (Mantel, 1907) The maximum number of edges in an n -vertex **triangle-free** graph is $\lfloor \frac{n^2}{4} \rfloor$.

Proof.

(i) *There is a triangle-free graph with $\lfloor \frac{n^2}{4} \rfloor$ edges.*

(ii) *If G is a triangle-free graph, then $e(G) \leq \lfloor \frac{n^2}{4} \rfloor$.*

Proof of (ii) is with extremality. (Look at the neighborhood of a vertex of maximum degree.)

Example of a wrong proof of (ii) by induction.

Bipartite subgraphs

Theorem. Every loopless multigraph G has a bipartite subgraph with at least $e(G)/2$ edges.

Proof # 1. Algorithmic. (Start from an arbitrary bipartition and move over a vertex whose degree in its own part is *more* than its degree in the other part. Iterate. Prove that at termination you have what you want.)

Proof # 2. Extremality. (Consider a bipartite subgraph H with the *maximum number of edges*, prove that $d_H(v) \geq d_G(v)/2$ for every vertex $v \in V(G)$ and use the Handshaking Lemma.)

Remark 1. *Maximum vs. maximal.* Algorithmic proof *not* necessarily ends up in bipartite subgraph with maximum number of edges.

Remark 2. The constant multiplier $\frac{1}{2}$ of $e(G)$ in the Theorem is **best possible**. *Example:* K_n .